

# Increased leisure-time physical activity associated with lower onset of diabetes in 44 828 adults with impaired fasting glucose

Lao, Xiang Qian; Deng, Han-Bing; Liu, Xudong; Chan, Ta-Chien; Zhang, Zilong; Chang, Ly-Yun; Yeoh, Eng-Kiong; Tam, Tony; Wong, Martin Chi Sang; Thomas, G Neil

DOI:

[10.1136/bjsports-2017-098199](https://doi.org/10.1136/bjsports-2017-098199)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Lao, XQ, Deng, H-B, Liu, X, Chan, T-C, Zhang, Z, Chang, L-Y, Yeoh, E-K, Tam, T, Wong, MCS & Thomas, GN 2018, 'Increased leisure-time physical activity associated with lower onset of diabetes in 44 828 adults with impaired fasting glucose: a population-based prospective cohort study', *British Journal of Sports Medicine*. <https://doi.org/10.1136/bjsports-2017-098199>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

Supplementary Table 14. Associations between volume of leisure-time physical activity and incident diabetes.

Coefficient between volume of leisure time physical activity (MET-hours/week) and incident type 2 diabetes)		
Model 1	Model 2	Model 3
-0.88 (-1.18 to -0.58) §	-0.66 (-1.04 to -0.28) ‡	-0.04 (-0.45 to 0.37)

Model 1: adjusted for age, sex;

Model 2: Model 1 + sex, marital status, education, physical labour at work, smoking, alcohol drinking, sleep duration, vegetable intake, systolic blood pressure, heart rate and total cholesterol;

Model 3: Model 2 + body mass index, waist circumference, fasting plasma glucose, triglycerides and white blood cell count.

‡ $P < 0.01$ ; § $P < 0.001$ .

Supplementary Table 2. Hazard ratios (95% CI) for incident diabetes by volume of leisure-time physical activity

	<u>Volume of leisure-time physical activity</u>				<u>Trend <i>P</i></u>
	<u>Inactive</u>	<u>Low</u>	<u>Moderate</u>	<u>High</u>	
<u>Model 1</u>	<u>1.00</u>	<u>0.90 (0.83-0.98) †</u>	<u>0.82 (0.74-0.90) §</u>	<u>0.74 (0.67-0.80) §</u>	<u>&lt; 0.001</u>
<u>Model 2</u>	<u>1.00</u>	<u>0.89 (0.80-0.98) †</u>	<u>0.80 (0.71-0.90) §</u>	<u>0.75 (0.67-0.83) §</u>	<u>&lt; 0.001</u>
<u>Model 3</u>	<u>1.00</u>	<u>0.96 (0.87-1.06)</u>	<u>0.93 (0.82-1.05)</u>	<u>0.89 (0.80-1.00)</u>	<u>0.034</u>

Inactive (no LTPA or LTPA < 3.75 MET-hours/week; reference category), low (LTPA 3.75 to < 7.5 MET-hours/week), moderate (LTPA 7.5 to < 15.0 MET-hours/week) and high (LTPA ≥ 15.0 MET-hours/week).

Model 1: adjusted for age, sex;

Model 2: Model 1 + marital status, education, physical labour at work, smoking, alcohol drinking, sleep duration, vegetable intake;

Model 3: Model 2 + body mass index, waist circumference, fasting plasma glucose, triglycerides and white blood cell count.

†*P* < 0.05; ‡*P* < 0.01; §*P* < 0.001.

Supplementary Table 43. Hazard ratios (95% CI) ~~of~~ for incident diabetes by volume of leisure-time physical activity with exclusion of incident diabetes identified within first 2 years.

	Volume of leisure-time physical activity				Trend <i>P</i>
	Inactive	Low	Moderate	High	
Case/Person-years	1780/103815	493/34424	364/21984	417/28841	
Model 1	1.00	0.88 (0.80-0.98) †	0.80 (0.72-0.90) §	0.72 (0.65-0.80) §	< 0.001
Model 2	1.00	0.87 (0.77-0.98) †	0.77 (0.67-0.90) ‡	0.72 (0.63-0.82) §	< 0.001
Model 3	1.00	0.94 (0.83-1.07)	0.89 (0.76-1.03)	0.86 (0.75-0.98)	0.013

Inactive (no LTPA or LTPA < 3.75 MET-hours/week; reference category), low (LTPA 3.75 to < 7.5 MET-hours/week), moderate (LTPA 7.5 to < 15.0 MET-hours/week) and high (LTPA ≥ 15.0 MET-hours/week).

Model 1: adjusted for age, sex;

Model 2: Model 1 + marital status, education, physical labour at work, smoking, alcohol drinking, sleep duration, vegetable intake, systolic blood pressure, heart rate and total cholesterol;

Model 3: Model 2 + body mass index, waist circumference, fasting plasma glucose, triglycerides and white blood cell count.

†*P* < 0.05; ‡*P* < 0.01; §*P* < 0.001.

Supplementary Table 24. Population attributable fractions (95% CI) ~~of~~ [for incident](#) diabetes by volume of leisure-time physical activity with exclusion of incident diabetes identified within first 2 years.

	Volume of leisure-time physical activity			
	Inactive	Inactive → Low	Inactive → Moderate	Inactive → High
No. participants	15439	5276	3303	4331
Case/Person-years	1780/103815	493/34424	364/21984	417/28841
Population attributable fraction (%)	----	10.56 (2.11-18.28)	18.62 (8.63-27.51)	15.36 (4.90-24.67)

[Inactive \(no LTPA or LTPA < 3.75 MET-hours/week; reference category\), low \(LTPA 3.75 to < 7.5 MET-hours/week\), moderate \(LTPA 7.5 to < 15.0 MET-hours/week\) and high \(LTPA ≥ 15.0 MET-hours/week\).](#) Population attributable fractions (PAFs) were calculated from hazard ratios that were adjusted for age, sex, marital status, education, physical labour at work, smoking, alcohol drinking, sleep duration, vegetable intake, systolic blood pressure, heart rate and total cholesterol.

Supplementary Table 35. Adjusted population attributable fractions (95% CI) ~~of~~ [for incident](#) diabetes by combinations of intensity and duration of leisure-time physical activity with exclusion of incident diabetes identified within first 2 years.

<b>Intensity</b>	<b>Duration</b>		
	Inactive → 0.5-2.4 hours/week	Inactive → 2.5 to 6.9 hours/week	Inactive → ≥ 7 hours/week
Inactive → Light (2.5 METs)	10.29 (-0.11-19.61)	13.24 (0.75-24.16)	17.05 (-7.02-35.70)
Inactive → Moderate (4.5 METs)	15.58 (-2.29-30.33)	19.41 (3.40-32.76)	50.6 (24.69-67.56)
Inactive → Vigorous (≥ 6.5 METs)	39.00 (19.65-53.68)	27.82 (11.26-41.29)	35.58 (7.75-55.01)

[Inactive \(no LTPA or LTPA < 3.75 MET-hours/week\)](#). Population attributable fractions (PAFs) were calculated from hazard ratios that were adjusted for age, sex, marital status, education, physical labour at work, smoking, alcohol drinking, sleep duration, vegetable intake, systolic blood pressure, heart rate and total cholesterol. The MET values for each intensity category were assigned by investigators according to the Ainsworth's compendium of physical activities [9].

Supplementary Figure 1. [Enrolment Selection](#) of [the](#) participants

